



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION OF PROB. 313 (SEE P. 164) BY PROF. D. V. WOOD. — There being no escape of heat due to compression, the law of pressure would be expressed by

$$pv^k = \text{constant} = p'v'^k, \quad (1)$$

where p' = initial pressure of the atmosphere = 15lbs, nearly, v' = vol. of the cylinder = al , if a is the area of the base, p = the pressure within the cylinder when the weight has descended a distance x , $a(l-x)$ = the volume when the pressure is p , $k = 1.408$; then, from (1),

$$p = \frac{p'l^k}{(l-x)^k}, \quad \text{and} \quad \frac{wd^2x}{gdt^2} = w + p'a - \frac{ap'l^k}{(l-x)^k}.$$

Integrating, observing that for $x = 0$, $v = 0$,

$$v^2 = \left(1 + \frac{p'a}{w}\right)2gx + \frac{2ap'g}{w(k-1)} \left(\frac{l[l-x]^{k-1} - l^k}{[l-x]^{k-1}} \right).$$

At the end of the downward movement $v = 0$; therefore

$$x(w + pa)(l-x)^{k-1} + \frac{ap'l}{k-1}(l-x)^{k-1} = \frac{ap'l^{k+1}}{k-1},$$

from which x may be found by trial after numerical values have been substituted for the known quantities.

SOLUTIONS OF PROBLEMS IN NUMBER FIVE.

SOLUTIONS of problems in No. 5, have been received as follows:

From Prof. W. P. Casey, 320, 321, 323; Prof. J. H. Kershner, 324 and Miscellaneous prob. No. 1 (see p. 149); Octavian L. Mathiot, 320; Prof. D. J. Mc Adam, 323; Prof. E. B. Seitz, 324; Prof. M. C. Woodward, 322; Prof. D. V. Wood, 322; H. Heaton, 322.

320. "The transverse and conjugate axes of an ellipse being given, to find the diameter of the circular base, and the altitude, of the right cone, and where to pass a plane so as to produce the given ellipse."

SOLUTION BY OCTAVIAN L. MATHIOT, BALTIMORE, MD.

Construction. — Draw $AB = 2a$, the given transverse diameter. At P , the centre of AB , erect a perpendicular, and with A as a centre and radius AB , describe a vertical arc in which suppose a point D to be taken such that if BD be produced to meet the perpendicular from P in C , and the triangle PBC be revolved about PC as an axis, a cone will be described a section of which, having AD for transverse axis, will have the given conjugate axis.